

The global biogeography of tree leaf form and habit

Received: 14 December 2022

Accepted: 18 September 2023

Published online: 23 October 2023

 Check for updates

A list of authors and their affiliations appears at the end of the paper

Understanding what controls global leaf type variation in trees is crucial for comprehending their role in terrestrial ecosystems, including carbon, water and nutrient dynamics. Yet our understanding of the factors influencing forest leaf types remains incomplete, leaving us uncertain about the global proportions of needle-leaved, broadleaved, evergreen and deciduous trees. To address these gaps, we conducted a global, ground-sourced assessment of forest leaf-type variation by integrating forest inventory data with comprehensive leaf form (broadleaf vs needle-leaf) and habit (evergreen vs deciduous) records. We found that global variation in leaf habit is primarily driven by isothermality and soil characteristics, while leaf form is predominantly driven by temperature. Given these relationships, we estimate that 38% of global tree individuals are needle-leaved evergreen, 29% are broadleaved evergreen, 27% are broadleaved deciduous and 5% are needle-leaved deciduous. The aboveground biomass distribution among these tree types is approximately 21% (126.4 Gt), 54% (335.7 Gt), 22% (136.2 Gt) and 3% (18.7 Gt), respectively. We further project that, depending on future emissions pathways, 17–34% of forested areas will experience climate conditions by the end of the century that currently support a different forest type, highlighting the intensification of climatic stress on existing forests. By quantifying the distribution of tree leaf types and their corresponding biomass, and identifying regions where climate change will exert greatest pressure on current leaf types, our results can help improve predictions of future terrestrial ecosystem functioning and carbon cycling.

Forest ecosystems, which contain 80–90% of global terrestrial plant biomass^{1,2} and a large proportion of terrestrial biodiversity³, regulate global biogeochemical cycles, and provide critical ecosystem services⁴. Leaves mediate forest energy and carbon inputs via photosynthesis, respiration, transpiration^{5,6} and litterfall^{7,8}, thereby regulating ecosystem structure and function, and water, nutrient and carbon cycles^{9–11}. Leaves of trees are highly diverse but can be broadly classified into four major types on the basis of leaf habit (evergreen vs deciduous) and form (broadleaved vs needle-leaved). These characteristics are linked to a vast array of functional traits associated with resource-use strategies and strongly depend on local growing conditions^{12–15}. Therefore, understanding variation in leaf types along

environmental gradients is critical to predicting global biogeochemical cycles and ecosystem functioning in a changing world. Yet, we still lack a global, quantitative understanding of forest leaf habit and form, informed by field-based observations.

Deciduous tree species evolved to tolerate seasonal climates and maximize the use of a short growing season^{16,17}. They usually have higher photosynthetic rates¹⁸ than evergreen species and reduce transpiratory water loss due to respiration by shedding their leaves during unfavourable seasons¹¹. Evergreen trees with longer leaf lifespans, by contrast, tend to have greater leaf construction costs¹⁹ and lower nutrient cycling rates²⁰. Growing season water-use strategy commonly differs between broadleaved and needle-leaved species²¹, with needle-leaved species

✉ e-mail: haozhi.ma@usys.ethz.ch




Fig. 1 | Global coverage of forest inventory locations (GFBi data) and plot-level leaf-type proportions. **a**, A total of 9,781 forest inventory plots (green points) were used for geospatial modelling of forest leaf types. **b**, Number of plots in relation to their proportion of evergreen vs deciduous and broadleaved vs needle-leaved individuals.

often showing conservative strategies²², such as lower stomatal conductance²³ and higher hydraulic safety margins²⁴, resulting in low photosynthesis rates²⁵. A spatially explicit understanding of tree leaf types is therefore critical for estimating the sensitivity and resilience of forests to future climate and soil conditions^{26–28}, and understanding the ecological consequences of such changes.

Theoretical models²⁹ and remote sensing observations^{30,31} have shown general trends in how climate and soil conditions affect the geographic occurrence of broadleaved and needle-leaved tree species at regional and global scales³². These relationships form the foundation of dynamic global vegetation models^{31,33–37}. Yet, the relative importance of various environmental characteristics on leaf habit and form remains to be determined. Furthermore, until now, these vegetation models have lacked the ground data needed to build tree-density-based ‘bottom-up’ models. Such models are crucial for validating satellite-derived trends on a global scale and for providing a comprehensive, high-resolution depiction of forest leaf-type variation across environmental gradients.

Here we analyse the global distribution of needle-leaved, broadleaved, evergreen and deciduous tree species, by integrating ground-sourced information from 9,781 standardized forest inventory plots in the Global Forest Biodiversity initiative (GFBi)³⁸ database with species-level leaf habit (evergreen vs deciduous) and leaf form (broadleaf vs needle-leaf) information accessed from the TRY plant trait database³⁹ (Fig. 1a). The 9,781 forest inventory plots represented a subsample of the full GFBi data of >1 million records to ensure an equal representation of different forest biomes across the globe and avoid modelling bias caused by uneven spatial sampling (see Methods). Using information on both the occurrences of individual trees per plot and the basal-area weighted occurrences of each individual, we calculated plot-level leaf-type proportions (1) on the basis of the leaf type of each individual tree (tree-based leaf type) and (2) by weighting each tree by its basal area (area-based leaf type) (see Methods section ‘Tree leaf-type data’). The first estimate allowed us to map the tree densities of each leaf type, while the second estimate allowed us to map the leaf types represented by the largest trees in a plot. To interpolate patterns across the globe, we combined our plot-level forest leaf information with 58 environmental variables, representing vegetation characteristics, climate, topography, vegetation, soil conditions and human-related features. We also tested the relative importance of 29 commonly studied variables on leaf-type variation and characterized the relationships between environmental features and leaf type.

Mapping global forest leaf types

To characterize the variation in forest leaf type, we first summarized the proportion of evergreen vs deciduous (leaf habit) and broadleaved vs needle-leaved (leaf form) individuals within each plot (Fig. 1a). Across all 9,781 forest plots, 13% contain exclusively broadleaved evergreen

trees, 7% contain exclusively broadleaved deciduous trees, 12% contain exclusively needle-leaved evergreen trees, and the remaining 68% contain a mixture of leaf habits and forms (Fig. 1b).

Our random forest model predicted 75% of the global variation of forest leaf-type classes (10-fold cross-validation R^2_{BC} , see Methods). We also ran spatially buffered leave-one-out cross-validation (LOO-CV) to account for the potential effect of spatial autocorrelation on model evaluation statistics, which resulted in a coefficient of determination (R^2) of 0.56 at a buffer radius of 300 km (see Methods and Supplementary Fig. 1). Within each class, our model explained 90%, 59%, 75% and 29% (10-fold cross-validation R^2) of the global variation in the proportion of broadleaved evergreen, broadleaved deciduous, needle-leaved evergreen and needle-leaved deciduous trees within forests, respectively. These predictive relationships were then used to upscale the observations across the global extent of forest coverage (Fig. 2).

To evaluate model robustness, we tested its performance on an independent validation dataset containing 3,895 sites across the globe⁴⁰, resulting in an R^2 of 0.47 (see Methods section ‘Cross-validation using external data’). In addition, we compared our model output with annual land cover maps from the European Space Agency’s Climate Change Initiative (ESA CCI LC)⁴¹. Across all leaf types, our model showed high correlations with the ESA CCI LC outputs, with an R^2_{BC} of 0.61. Within each leaf type class, our model explained 78% of the variation in the proportion of broadleaved evergreen trees, 31% of broadleaved deciduous tree proportions, 64% of needle-leaved evergreen tree proportions and 19% of needle-leaved deciduous tree proportions (R^2).

While considerable uncertainties exist for individual predictions at the pixel level, these uncertainties rapidly decrease as the model is projected to a larger area (<5% at 250 km²; Supplementary Fig. 2). Our model shows high confidence in tropical and boreal forests, whereas predictive confidence is lower in mixed forests and ecotones between different types of forest (Supplementary Figs. 3 and 4). For example, models for broadleaved evergreen and deciduous species share low predictive confidence in central African savanna regions. Similarly, low predictive confidence can be found across eastern Russian mixed forests. The low predictive confidence for these regions can be attributed to low sample size and mixed occurrence of broadleaved evergreen and deciduous species, as well as differences in the year of observation across forest survey plots, which may lead to larger uncertainties in ecotones where forest types can shift in relatively short time periods.

Global environmental drivers of forest leaf-type variation

To assess the relative importance of environmental features on variation in leaf types, we ran random forest models including a range of environmental variables (see Methods). We combined these environmental factors into three groups (climate, soil and topography).




Fig. 2 | The global distribution of forest leaf types. **a**, The global distribution of tree leaf type as predicted by a random forest model built from area-based leaf-type data (see Methods). Pixels are coloured in the red, green and blue spectrum according to the percentage of total tree basal area occupied by broadleaved evergreen, broadleaved deciduous and needle-leaved tree types, as indicated by the ternary plot. Needle-leaved evergreen and needle-leaved deciduous forests

were combined due to the low global coverage of needle-leaved deciduous trees. **b–e**, Predicted relative coverage of each leaf type from random forest models. Ref. [81](#) was used to mask non-forest areas. **b**, Broadleaved evergreen coverage. **c**, Broadleaved deciduous coverage. **d**, Needle-leaved evergreen coverage. **e**, Needle-leaved deciduous coverage.

To test for the relative importance of climate, soil and topographic characteristics, we ran a principal component analysis (PCA) for each of these variable groups and selected the first six principal components, which explained $\geq 90\%$ of the total variation across all variables. Our analyses show that climate and soil characteristics jointly determine the global leaf-type distribution (Fig. [3a](#)). With respect to variation in leaf habit, temperature fluctuations, that is, isothermality and temperature seasonality, were the most important variables (Fig. [3c](#)). Yet, the entire combination of soil features (first six principal components of soil variables) was as important as climate for predicting leaf habit in our random forest model (Fig. [3a](#)), suggesting that soil characteristics play an important role in the global distribution of evergreen ‘vs’ deciduous species. Especially soil texture, in combination with pH, appears to affect global variation in tree leaf habit. Acidic soils, commonly found in tropical and boreal regions, inhibit nutrient (N and P) supply by reducing cation availability and limiting tree

growth rates¹⁶. This might explain why broadleaved deciduous species that require high nutrient supply are less abundant in regions with acidic soil. Broadleaved evergreen species, by contrast, may better cope with nutrient poor, acidic soils⁴². Similarly, needle-leaved evergreen species that can maintain growth even under low nutrient supply are more abundant in regions with acidic soil¹⁶. The high tannin and phenol contents of needle leaves further contribute to the acidification of soils¹⁶, probably creating a positive feedback towards coniferous dominance. Overall, our results point towards a feedback between tree leaf habits and soil conditions, highlighting the connection between physical soil features and soil water⁴³ and nutrient⁴⁴ availability.

Variation in leaf form was best predicted by climate variables (Fig. [3b](#)), with the most important variable being temperature of the coldest quarter (Fig. [3d](#)). By contrast to leaf habit, soil, topographic and vegetation features were less important in driving variation in leaf form, indicating adaptation to extreme climates, cold winters or




Fig. 3 | Variable importance of environmental covariates on forest leaf-type variation. **a,b**, Cumulative importance of the first six principal components of climate, soil, topographic and vegetation covariates in the variation of leaf habit (**a**) and leaf form (**b**). **c,d**, Variable importance of selected environmental features

on variation in leaf habit (**c**) and leaf form (**d**). Bars in **c** and **d** represent the mean $\pm$ 95% CI; relative importance based on the 10 best random forest models ($n = 10$; see Methods). Area-based leaf-type proportions were used to represent forest (plot-level) leaf-type variation.

extended dry seasons, as a major determinant of leaf form. This supports previous studies indicating that diverse leaf forms evolved to adapt to different climates^{45,46}.

Computation of tree densities with different leaf types

To quantify the proportions of different leaf types across global forests, we combined a global tree density distribution map⁴⁷ with our individual-based leaf type models (see Methods and Supplementary Fig. 5). At the global scale, we estimate that of the ~3 trillion adult trees presently existing, 29.1% (95% CI = 27.5–30.7%) are broadleaved evergreen, 27.1% (23.8–30.6%) are broadleaved deciduous, 38.4% (35.2–41.6%) are needle-leaved evergreen and 5.4% (4.3–6.6%) are needle-leaved deciduous (Fig. 4 and Supplementary Fig. 6). Even though needle-leaved tree species comprise less than 2% of the world's estimated 73,000 tree species⁴⁸, this small fraction of diversity represents around 44% of individual trees on Earth.

Of the 1.15 trillion needle-leaved evergreen trees growing worldwide, the majority (64.7%) are found in boreal regions, while 23.1%, 8.9%, 2.8% are found in temperate, arid and tropical regions, respectively (Fig. 4). In contrast, of the 0.87 trillion broadleaved evergreen trees growing worldwide, the majority exists in tropical (57.2%) and arid (29.8%) regions, with 6.3% and 4.3% existing in temperate and boreal regions, respectively (Fig. 4). Broadleaved deciduous trees show the widest range of occurrences. Of the 0.81 trillion broadleaved deciduous trees, 34.5% are found in boreal regions, 28.8% in temperate regions,

22.5% in arid regions and 11.5% in tropical regions (Fig. 4). We further estimate that there are 0.16 trillion needle-leaved deciduous trees across the globe, the vast majority of which grow in boreal regions.

Using our basal-area-based model of leaf types, we were able to estimate the biomass contribution of each leaf type within individual forest pixels by integrating our data with a recently published aboveground biomass map⁴⁹. Our analysis revealed that broadleaved evergreen trees store the largest proportion of global forest biomass, accounting for 54.4% (335.7 Gt) out of the total biomass of 617 Gt. Broadleaved deciduous trees contribute 22.1% (136.2 Gt), needle-leaved evergreen trees contribute 20.5% (126.4 Gt) and needle-leaved deciduous trees contribute 4% (18.7 Gt) (Supplementary Fig. 7). Interestingly, despite there being 42% more evergreen needle-leaved trees compared with broadleaved evergreen trees, their biomass contribution is 62% (209.3 Gt) lower. This distribution of biomass across different leaf types provides valuable insights into the carbon storage capacity of diverse forest ecosystems.

Climatic risk assessment of future leaf types

Climate change will strongly affect the functioning of terrestrial ecosystems by altering growth, mortality and reproduction of trees and their interactions with leaf form and habit¹⁶. Our models allowed us to highlight areas of potential risk by identifying the regions where future climates will shift to conditions that currently support leaf types different from the prevailing ones. In these regions, trees are likely to experience more climatic stress in the future. To assess the




Fig. 4 | The global proportion of evergreen broadleaved, deciduous broadleaved, needle-leaved evergreen and needle-leaved deciduous trees. The relative proportions of trees that occur within tropical, temperate, boreal and arid regions are shown as separate pie charts for each leaf type.

extent and distribution of future changes in forest leaf-type climate envelopes, we projected our leaf-type models into the future using three climate change scenarios (low-emission scenario (SSP1–RCP2.6), business-as-usual scenario (SSP3–RCP7) and high-emission scenario (SSP5–RCP8.5))⁵⁰. To do so, we used our random forest models of present leaf type distributions and replaced all climate variables reflecting the 1979–2013 climate (see Supplementary Fig. 1) with climate model projections for 2071–2100 while keeping soil, topographic, vegetative and anthropogenic characteristics constant.

The results suggest that forests will experience substantial shifts in leaf-type climate conditions. Depending on future emissions pathways, 17 to 34% of future forested regions are projected to experience a climate by the end of the century that currently supports leaf types different from the prevailing ones (Fig. 5 and Supplementary Fig. 8; see Supplementary Fig. 9 for an alternative definition of forest types). The climate conditions that have historically supported evergreen forests are declining as global conditions shift towards those that have historically supported more deciduous forests, and this appears to be the case for both broadleaved and needle-leaved species (Supplementary Fig. 10). Specifically, 7–20% of the broadleaved evergreen forests are likely to experience changes towards conditions that currently support deciduous species (Fig. 5). Similarly, 29–67% of the needle-leaved evergreen forests will experience changes towards climate conditions that currently support mixed or deciduous forests (Fig. 5 and Supplementary Figs. 8–12). If these climate projections are realized, plants in those regions must either tolerate more stressful environmental conditions or shift their distributions, causing changes in forest composition⁵¹. Previous studies predicting ecoregion shifts have also shown a heightened vulnerability to changing climate conditions, surpassing even the susceptibility of leaf types⁵².

It is important to acknowledge that linking forest leaf types to climate alone cannot fully capture the complex interactions of

other influential factors, including CO₂ concentration^{53,54} and nutrient availability²⁶. Consequently, the analysis presented in this study does not project actual changes in forest leaf types. Instead, its focus is on identifying regions where future climates will shift to conditions that currently support different leaf types than what is currently observed. Different species may exhibit varying tolerance thresholds and responses to CO₂ fertilization, leading to divergent outcomes. Moreover, it is possible that species sharing the same leaf habit or form but with broader climatic tolerance ranges could replace the present species, potentially mitigating leaf-type changes.

Our analysis serves as a risk assessment, highlighting regions where climate poses a potential threat to the existing forest composition. Further research is necessary to gain a deeper understanding of the intricate interactions between climatic changes, elevated atmospheric CO₂ concentrations and nutrient availability. These interactions play a pivotal role in determining essential aspects of forest ecology, such as germination rates, seedling survival, growth rates and tree mortality, which ultimately shape forest composition. Nonetheless, the analysis underscores the substantial and rapid changes in climatic conditions that forests are already experiencing and will continue to experience even more profoundly in the future, on timescales ranging from decades to centuries.

Methodological considerations

Our models successfully captured a substantial portion of the observed spatial variation in forest leaf types, with an overall explained variation of 75%. However, the accuracy of our predictions varied across different leaf types. Among them, our model achieved the highest accuracy in predicting the spatial distribution of broadleaved evergreen tree species, with an explained variation of 90%. Conversely, our model explained only 29% of the variation in needle-leaved deciduous trees, which can be attributed to the limited availability of data from plots containing needle-leaved deciduous species.




Fig. 5 | Forested areas where future climates may no longer support prevailing leaf types. If a pixel's forest area was predominantly (>60%) covered by one leaf type, it was classified as that specific leaf type. Pixels where no leaf type exceeded 60% coverage were classified as mixed forest. To determine the relative proportion of each leaf type per plot, we considered the basal area of individual trees (area-based leaf type). Coloured pixels on the map indicate areas that, by the end of the century (2071–2100), will face climate conditions that

currently support a different forest type. The future climate conditions were represented using three climate change scenarios: low-emission (SSP1–RCP2.6; **a,b**), business-as-usual (SSP3–RCP7; **c,d**) and high-emission (SSP5–RCP8.5; **e,f**) for the period 2071–2100. Panels **a, c** and **e** show the present forest types. In contrast, panels **b, d** and **f** show the type of forest expected under the projected future climate of each respective pixel.

To address the uncertainties associated with our predictions, we employed a subsampling approach, running 100 independent models. This allowed us to assess the range of variability in our predictions and identify areas with lower predictive confidence. The resulting maps of model uncertainty (Supplementary Fig. 3) highlight regions where caution should be exercised when interpreting our predictions.

While our study supports many of the mechanisms identified in previous research, our correlative analyses do not necessarily establish causal relationships. To map leaf-type variation across the globe on the basis of the relationships with environmental features, we used global layers. This approach was necessary as point-level predictor variables cannot be used to interpolate predictions across the globe. While the majority of these layers effectively capture local variations, soil layers may not fully reflect point-level soil observations due to the inherent spatial heterogeneity of soil conditions. In addition, soil layers are typically derived from interpolation methods using environmental information and may thus be correlated with climate variables. While our random forest model predictions are not affected by multicollinearity, this could impact the quantification of variable importance related to environmental, soil, topographic and vegetation features.

Incorporating local point observations into the model training was not feasible because the ground-based forest inventory data we used did not include field-measured environmental and soil characteristics.

Although we examined additional point-level observations from the World Soil Information Service (WOSIS) dataset (see next paragraph), these observations often did not align spatially with the forest inventory plots, resulting in a reduced sample size (>80% reduction) and geographic bias (Fig. 1 and Supplementary Fig. 14). Moreover, training the model using point observations while relying on soil layers for model prediction could introduce further uncertainties and biases. To avoid these limitations, we conducted both model training and predictions using the same soil layers.

Nevertheless, we conducted analyses to determine the agreement between results based on global soil layers and point-level soil observations. We matched our forest inventory dataset⁵⁵ with the WOSIS dataset⁵⁶, which contains local point observations of soil features. These analyses indicated that: (1) model predictions remained consistent when using point observations instead of global layers of soil characteristics (97–99% similarity in predictions), (2) the global soil layers exhibited good agreement with point observations for most soil variables, particularly the four most important variables ($R^2 = 0.42–0.62$; Fig. 3c and Supplementary Fig. 13) and (3) the inferred importance of soil features in leaf type variation remained similar (<5% difference) when using point observations instead of soil layers as predictors (Supplementary Fig. 14). These analyses underscore the crucial role of soil features in determining global leaf-type variation.

Conclusions

By characterizing the environmental controls of forest leaf-type variation and integrating this information with tree density data, our analysis reveals that 38% of global tree individuals are needle-leaved evergreen, 29% are broadleaved evergreen, 27% are broadleaved deciduous and 5% are needle-leaved deciduous. We further show predictable global patterns in forest leaf varieties, with the relative abundance of leaf forms mainly correlating with temperature variation. In contrast, leaf habit is jointly controlled by climate and soil characteristics. These global relationships between environmental factors and forest leaf types largely agree with local experimental and modelling studies^{57–59}, which also highlight the dual role of climate and soil conditions in driving variation in forest types.

Our analysis of the spatial variation of forest leaf types refines our understanding of forest composition and structure^{60,61} at a global scale. While satellite-derived approaches have been foundational for the characterization of spatial variation in canopy structure⁶², our bottom-up model, derived from empirical inventory data, allowed us to create models of forest leaf type at the individual tree level to provide novel insights into forest composition. This can help benchmark satellite-derived models of forest structure and inform ecological models of plant productivity, biogeochemical cycling, carbon storage and species distribution^{31,63,64}. By identifying the main environmental characteristics that determine habitat suitability, such as annual mean temperature, climate seasonality and water and nutrient supplies, our baseline estimates of leaf-type densities are also critical for projecting population- and community-level tree demographics under current and future climate change. Ultimately, these insights can help make informed decisions to guide global efforts to conserve, restore and sustainably manage forest ecosystems that are so vital for the wellbeing of all organisms on Earth.

Methods

Data collection

Forest inventory data. To obtain empirical information on tree occurrences, we extracted a total of 1.1 million forest inventory plots with more than 50 million individual occurrence records, covering all continents except Antarctica from the GFBi dataset³⁸. To avoid modelling bias caused by uneven spatial sampling across biomes, we downsampled the dataset so that the relative proportion of plots in each biome in the dataset approximately matched the proportion of forested area within each biome⁶⁵. This was done by retaining all tropical forest plots in GFBi ($n = 11,367$) and randomly downsampling the remaining biomes in proportion to this number. Individuals with stem diameters <10 cm were excluded as the focus was on adult trees, and only plots with ≥ 10 adult individuals were included in the final analysis. For each plot, the dataset contains information on the location (coordinates), the year(s) when the inventory took place, stem diameter at breast height (DBH) and species identity of each individual. For plots with time series data, only the most recent observation year was included in the analysis. The average year of observation across all plots was 2005. This resulted in a total of 9,781 plots with a median size of $\sim 500 \text{ m}^2$ and 817,091 individual tree records, with 20.3% of the plots in the boreal biome (vs 21% of forested land globally), 20.4% in temperate biomes (vs 22%), 54% in tropical biomes (vs 50%) and 3% in Mediterranean woodland, tundra, xeric shrubland or mangrove biomes (vs 7% globally).

Tree leaf-type data. Information on leaf habit (evergreen vs deciduous) and leaf type (broadleaved vs needle-leaved) was extracted from the TRY plant trait database³⁹. For each species and genus, we assigned the most common leaf habit across all TRY observations, treating leaf habit as a binary variable of whether it is evergreen or not. Species names were standardized using the Taxonstand R package⁶⁶. We then assigned leaf-type information to the individuals included in the GFBi dataset using species-level information or genus-level information in

case species-level information was not available. Plots in which <50% of all individuals had a leaf-type record in TRY were excluded. Out of 10,274 species recorded globally in the GFBi dataset, leaf-type records could be assigned to 8,642 tropical tree species, 453 temperate tree species, 46 boreal tree species and 1,124 tree species in dry areas. Next, we calculated the proportion of each leaf-type combination (evergreen broadleaved, deciduous broadleaved, needle-leaved evergreen or needle-leaved deciduous) within a plot, either by dividing the sum of individuals featuring the respective leaf type by the total number of individuals within the plot (individual-based leaf type) or by dividing the basal area featuring the respective leaf type by the total basal area of all trees in the plot (area-based leaf type).

Environmental covariates. In total, 58 global environmental layers reflecting climate⁵⁰, topography, vegetation, anthropogenic and soil⁶⁷ (at 0 cm to 100 cm depth) characteristics were used as covariates in our analyses (Supplementary Fig. 1). Climate variables reflect the average climate between 1979–2013. All layers were standardized to 30-arc-second resolution ($\sim 1 \text{ km}^2$ at the equator).

Geospatial modelling

Random forest modelling. To predict the occurrence probabilities of the four forest leaf types (broadleaved evergreen, broadleaved deciduous, needle-leaved evergreen and needle-leaved deciduous), we ran random forest models in Google Earth Engine⁶⁸. In random forest, unlike traditional regression, correlation among variables does not affect model accuracy. Indeed, the ability to use many correlated predictors is one of the key benefits of machine learning models⁶⁹. When variables are correlated, the effect of these variables is ‘shared’ across the trees in the random forest. Because random forest does not estimate coefficients as in regression, this correlation does not hinder model fit or performance, but rather complicates efforts to quantify variable importance, which is also shared across correlated variables. Thus, by including numerous variables, even if correlated, we can improve our predictive power of the model to accurately quantify current carbon.

To run the models, we set the output mode to ‘MULTIPROBABILITY’ and randomly sampled 10 individuals from each plot 100 times, weighting individuals by their basal area to model area-based leaf types. The following equation was used to transform DBH to basal area:

$$A = \frac{\pi \text{DBH}^2}{4}$$

To model individual-based leaf types, individuals were sampled without weighting them. This resulted in 100 training datasets, each containing 98,330 rows. After extracting pixel values from 58 environmental covariates, we then modelled leaf types for each training dataset with a random forest model, using the 58 environmental covariates. The correlation between our point-level response variable (leaf type) and spatially contiguous covariates then allowed us to map the global distribution of leaf types.

To train global models of forest leaf types, we first ran a grid-search procedure, exploring the results of a suite of random forest models with different hyperparameters. The hyperparameters that were allowed to vary were the number of random trees (10, 20, 50, 100 and 250), the number of variables sampled at each split (1, 2, 4, 5, 8, 10, 15, 20 and 30) and the minimum sample size at the end of the nodes (1, 2, 5, 10, 15, 20 and 30); subsampling rate was constant at 0.632. To quantify predictive accuracy, we used the Bhattacharyya coefficient to compare the predicted and observed probabilities for each pixel, as is commonly used in image processing and feature extraction^{70–73}. Because four probability classes within a pixel are not independent, we cannot use standard loss functions to estimate predictive accuracy. Instead, we used the Bhattacharyya coefficient, given by $\sum_{i=0}^n \sqrt{O_i \times P_i}$, which quantifies the similarity between two vectors, O and P , with n categories. The Bhattacharyya coefficient falls within (0,1), equaling one only when $P_i = O_i$ for all i within a pixel (that is, when the

predictions exactly match the observed) and zero when the vectors are completely disjoint. To evaluate overall model performance, we then adopted a similar approach to that of ref. 74 for multinomial data⁷⁵, using the Bhattacharyya coefficients to calculate a pseudo- R^2 (R_{BC}^2) (equations 1–5) on the basis of 10-fold cross-validation:

$$R_{BC}^2 = 1 - \frac{MAE_{\text{model}}}{MAE_{\text{mean}}} \quad (1)$$

in which

$$MAE_{\text{model}} = \frac{\sum E_i}{N} \quad (2)$$

$$E_i = 1 - \sum_{i=1}^n \sqrt{O_i \times P_i} \quad (3)$$

and

$$MAE_{\text{mean}} = \frac{\sum E_{i\text{mean}}}{N} \quad (4)$$

$$E_{i\text{mean}} = 1 - \sum_{i=1}^n \sqrt{O_i \times \bar{O}_i} \quad (5)$$

where, MAE is the mean absolute error, O_i is the observed coverage of leaf type i , P_i is the predicted coverage of leaf type i (on out-of-fit data via cross-validation), $\bar{O}_i$ is the average coverage of leaf type i across all the observations and n is the number of forest leaf types (here, $n = 4$). Note that the summation terms in equations (3) and (5) are the Bhattacharyya coefficients between the observed multinomial distribution and the predicted distribution (equation 3) and the average distribution (equation 5). Thus, equation (3) is the predictive loss term for each pixel, with the MAE_{model} in equation (2) giving the average across all pixels, which equals zero only when the predictions perfectly match the observations. Similarly, equation (5) is the loss term for each pixel when using group-level means, with MAE_{mean} in equation (4) giving the average loss across all pixels. By comparing MAE_{model} to MAE_{mean} , we follow the standard approach for computing R^2 by quantifying performances relative to human predictions, with $R^2 = 1 - MAE_{\text{model}}/MAE_{\text{mean}}$ equalling 1 only when our predictions are perfect ($MAE_{\text{model}} = 0$) and R^2 being ≤ 0 when our predictions are equal to or worse than the mean. Importantly, as suggested in ref. 74, equation (1) is estimated using out-of-fit cross-validated data, where the predicted values are estimated by omitting the corresponding observed values from the training data, with the resulting pseudo- R^2 used to assess our four-element model output.

To create the final maps (Fig. 2 and Supplementary Fig. 5), we used the random forest model for each training dataset with the optimal suite of hyperparameters based on the R_{BC}^2 from the grid search. Extrapolation of our predictions across global forest areas resulted in 100 four-band global layers, with each band representing the global probability of one forest leaf type. We averaged the predictions from these 100 model outputs by taking the mean for the final map. We calculated the 95% confidence intervals across the 100 model layers⁷⁶ to represent sampling uncertainty.

As an alternative mapping approach, we used an independent tree-based classification and regression trees (CART) model⁷⁷ (Supplementary Fig. 15). This approach was used to test whether model performance depends on model type. If the two models (random forest and CART) show similar results, this indicates that predictions are not biased by model selection. Using the same independent ‘Tallo’ dataset⁴⁰ used for testing the robustness of the random forest model, the CART model had an explanatory power of 0.46, which is similar to the R_{BC}^2 of 0.47 of the random forest model. When directly comparing both models, the CART model showed 87% similarity (R_{BC}^2) with

the random forest model. This suggests that our maps and predictions do not depend on the type of model, and we report the random forest model results throughout the main text since it showed slightly higher accuracy⁵⁵.

Cross-validation using external data. We tested for the performance and correlation between the predictions of the area-based and individual-based random forest models. When using the same independent ‘Tallo’ dataset⁴⁰ for testing the robustness of the random forest models, the tree-based and area-based models had an explanatory power of 46% and 47%, respectively. When directly comparing both models, the area-based model showed 89% similarity (R_{BC}^2) with the individual-based model, showing that both metrics result in similar predictions of leaf-type variation.

To further evaluate the performance of our models, we additionally compared the model predictions with satellite-derived leaf type estimates from annual land cover maps from the ESA CCI LC⁴¹. We used land cover layers for the years 2000, 2005, 2010 and 2015, each with a spatial resolution of 300 m $\times$ 300 m. To assign each pixel to forest leaf-type classes, we recalculated the leaf-type proportions for each layer as these represented leaf-type proportions across all ecosystem types, including grasslands. For example, we recalculated forest leaf-type proportions for pixels with 30% broadleaf deciduous forest cover, 20% needle-leaf evergreen forest cover, 10% needle-leaf deciduous forest cover and 40% non-forest cover by dividing each forest cover percentage by the total area covered by forest, resulting in 50% broadleaved deciduous, 33.3% needle-leaved evergreen and 16.7% needle-leaved deciduous. We then calculated the average values across the four years for each pixel and compared the results with our model outputs. Our area-based models explained 61% of the spatial variation in the ESA CCI LC models, with an explanatory power of 78% for broadleaved evergreen leaf-type proportions, 31% for broadleaved deciduous, 64% for needle-leaved evergreen and 19% for needle-leaved deciduous.

To generate global layers of soil features, the Soil Grids dataset relies on machine learning models informed by global, spatially explicit information on various climate variables. This global interpolation of soil information using climate data may thus lead to an over-estimation of the covariation between climate and soil layers while reducing small-scale heterogeneity in soil features. To assess whether this potential caveat affects our results, we used point-level soil measurements from the WOSIS dataset, including clay content, silt content, pH and sand content. To spatially match this dataset with the full GFBi dataset containing more than 1.1 million plots across the globe⁵⁵, we used the ‘geosphere’⁷⁸ R package. We then selected the nearest soil observation that fell within 250 m or 1,000 m of the centre of each forest plot. This resulted in a spatial match between soil measurements and forest plots in 146 cases for the 250 m radius and in 1,893 cases for the 1,000 m radius (Supplementary Fig. 14a). To test whether model performance and predictions change when using point observations of soil features instead of global layers, we then trained random forest models using either WOSIS or Soil Grids soil data along with information on climate, topography, human and vegetation characteristics (Supplementary Fig. 14b–d). For both the 250 m and 1,000 m buffer radii, the models showed a high degree of agreement ($R_{BC}^2 = 0.99$) between model predictions.

In a second step, we then tested whether the use of point observations vs global layers of soil features affects the estimated importance of variables in driving leaf-type variation. The analysis showed a slight reduction in the importance of soil variables of 5–6% when using point observations rather than Soil Grids data (Supplementary Fig. 13), which is probably driven by the slightly lower covariation of soil point data with climate variables (Supplementary Fig. 16). Nevertheless, the results remain similar, showing that this difference is unlikely to affect the conclusions of our study.

Interpolation vs extrapolation in model predictions. To evaluate how well our training dataset represents the full multivariate environmental covariate space, we performed a principal-component-analysis-based approach following refs. 76,79. We projected the covariate composite into the same space using the centreing values, scaling values and eigenvectors from the principal component analysis of the training data. We created the convex hulls for each of the bivariate combinations from the top principal components and classified whether each pixel falls in or outside each of these convex hulls. We used 24 principal components with 276 combinations for all covariates for the sampling dataset. This analysis showed that 99.2% of land pixels (778,975,911 of 785,150,461) cover at least 90% of the environmental variables present in our training data locations (Supplementary Fig. 17).

LOO-CV. To account for the potential effect of spatial autocorrelation in model residuals on model validation statistics, we ran spatially buffered LOO-CV for a series of buffer radii from 10 m to 500 km. In LOO-CV, each observation is predicted on the basis of a model that includes all data outside the respective buffer radius. This results in 9,781 (=total number of observations) separate models for each buffer radius. Model performance was evaluated on the basis of R^2_{BC} . To assess the range of spatial autocorrelation, we ran semi-variograms for random cross-validation and LOO-CV model residuals in each forest type, showing that regardless of buffer radius or validation type, our residuals show weak spatial autocorrelation (Supplementary Fig. 1). Nevertheless, when eliminating any potential effects of spatial autocorrelation on model performance by applying large buffer radii of 300 km and 500 km, the out-of-sample R^2_{BC} remained high (0.56 and 0.53, respectively).

Global tree density and biomass calculation of leaf types

Tree leaf-type densities were estimated by integrating a map of the global tree density distribution⁴⁷ with our individual-based forest leaf-type maps. For each pixel, we multiplied tree density values with modelled forest leaf-type proportions to obtain the pixel-level stem density of each leaf type. We then summed up these pixel-level abundances across the globe to estimate the total abundances of each forest leaf type. To obtain the total number of trees of each leaf type for the major forest types, we defined tropical forests as pixels falling in the biomes tropical and subtropical moist broadleaf forest (WWF⁸⁰ biome 1), tropical and subtropical coniferous forest (biome 3) and mangroves (biome 14). Temperate forests were defined as pixels in the biomes temperate broadleaf and mixed forest (biome 4) and temperate coniferous forest (biome 5). Boreal forests were defined as pixels in the biomes boreal forest or taiga (biome 6), montane grasslands and shrublands (biome 10) and tundra (biome 11). Dry forests were defined as pixels in the biomes tropical and subtropical dry broadleaf forest (biome 2), tropical and subtropical grasslands, savannas and shrubland (biome 7), temperate grasslands, savannas and shrubland (biome 8), Mediterranean forests, woodlands and scrub (biome 12), and deserts and xeric shrubland (biome 13).

Forest biomass for each leaf type was computed by incorporating a map of global forest biomass⁴⁹ with our area-based leaf-type models. We calculated the absolute biomass by scaling biomass density with tree canopy cover⁸¹ and pixel area within each pixel. This absolute biomass per pixel was then partitioned by leaf-type proportions, derived from our area-based models. By summing up the pixel-level biomass across the globe, we were able to approximate the total amount of biomass stored in each of the leaf types.

Environmental drivers of forest leaf-type variations

To evaluate the relative importance of environmental features on forest leaf-type variation, out of the total 58 environmental covariates, we tested the effects of 29 commonly used environmental characteristics using random forest models (see Supplementary Fig. 1). We separated

the environmental features into four major groups: climate, soil, topography and vegetation. To equally represent each of the three groups in the model and minimize collinearity, we selected the first six principal components from climate, soil and topography groups. These six principal components explained $\geq 90\%$ of the total variation across all group variables. We included all the three vegetation characteristics, which are forest age, tree density and canopy height, into the analysis without computing the principal components. We then computed the variance inflation factors (VIFs) across all 21 principal components using the R package HH⁸². All VIFs were lower than 4, suggesting sufficient independence among covariates. The principal components were then used as predictors in random forest models with different combinations of hyperparameters (that is, 1 to 12 samples per split and a minimum sample size at the end of the nodes of 1 to 10), and we selected the ten best models with the highest coefficient of determining variation (R^2). Variable importance was determined by calculating the relative influence of each variable, expressed by the variable's attributed reduction in squared error. To quantify the importance of individual environmental factors, we used the same combinations of hyperparameters. Based on R^2 , the ten best random forest models were again chosen to explore the relative importance of each factor. The random forest models were run via the R package h2o⁸³.

Forest types and their future climate envelopes

To assess the extent and distribution of future changes in forest leaf-type climate envelopes, we projected our leaf-type models into the future using CMIP6 climate models for the time interval of 2071–2100 and three emission scenarios (SSP1–RCP2.6, SSP3–RCP7 and SSP5–RCP8.5)⁵⁰. The CMIP6 climate models were extracted following the ISIMIP3b protocol and included five models (gfdl-esm4, ukesm1-0-11, mpi-esm1-2-hr, ipsl-cm6a-lr and mri-esm2-0). We projected the global forest leaf-type distribution for each emission scenario on the basis of each of the five climate models using our random forest models. To do so, we used our 100 random forest models of present leaf-type distributions and replaced the climate variables ('bioclim' variables from the CHELSA dataset, marked with hashtags in Supplementary Fig. 1) with CMIP6 climate model projections while keeping soil, topographic, vegetative and anthropogenic characteristics constant. For each emission scenario and CMIP6 model, we aggregated the 100 layers by taking the mean. We then aggregated the five CMIP6 model projections by taking the mean. For both evergreen vs deciduous and broadleaf vs needle-leaf proportions, we then subtracted the present predictions by the averaged model projections for the three climate change scenarios. We then summarized the predictions for each of the three climate change scenarios across latitude, aggregating predictions for each half degree. This allowed us to identify areas where future climates will shift to conditions that currently support leaf types different from the prevailing ones (Fig. 5 and Supplementary Fig. 9). To do so, we first obtained information on the forest type that currently is most abundant in each pixel (using area-based leaf-type proportions). Pixels in which $> 60\%$ of the forest area was covered by a single leaf type were assigned to that respective leaf type. Pixels in which none of the leaf types covered $> 60\%$ of the forest area were categorized as mixed forest (Fig. 5). The analysis was also conducted for a forest-area threshold of 80% to ensure that the results are not driven by the choice of the threshold (Supplementary Fig. 9). Following these definitions, we categorized forest pixels into groups using present and future climate scenarios. By scaling pixels by canopy cover, we could then calculate the total areas in which the climate is expected to shift to conditions that currently support a different forest type.

Reporting summary

Further information on research design is available in the Nature Portfolio Reporting Summary linked to this article.

Data availability

Tree occurrence data from the Global Forest Biodiversity initiative (GFBi) is available upon request via Science-I (<https://science-i.org>) or the GFBi website (<https://www.gfbiiinitiative.org/>). Information on leaf habit (evergreen vs deciduous) and leaf form (broadleaved vs needle-leaved) came from the TRY database (<https://www.try-db.org>). Additional, leaf-type data came from the Tallo dataset (<https://zenodo.org/record/6637599>). Plot-level soil information came from the World Soil Information Service (WOSIS) dataset (<https://www.isric.org/explore/wosis>).

Code availability

All code is available at <https://doi.org/10.5281/zenodo.7967245>.

References

- Pan, Y., Birdsey, R. A., Phillips, O. L. & Jackson, R. B. The structure, distribution, and biomass of the world's forests. *Annu. Rev. Ecol. Syst.* **44**, 593–622 (2013).
- Pan, Y. et al. A large and persistent carbon sink in the world's forests. *Science* **333**, 988–993 (2011).
- The State of the World's Forests 2020. Forests, Biodiversity and People* (FAO and UNEP, 2020).
- Bonan, G. B. Forests and climate change: forcings, feedbacks, and the climate benefits of forests. *Science* **320**, 1444–1449 (2008).
- Wright, I. J. et al. The worldwide leaf economics spectrum. *Nature* **428**, 821–827 (2004).
- Schulze, E. D. Biological control of the terrestrial carbon sink. *Biogeosciences* **3**, 147–166 (2006).
- Sayer, E. J. Using experimental manipulation to assess the roles of leaf litter in the functioning of forest ecosystems. *Biol. Rev. Camb. Phil. Soc.* **81**, 1–31 (2006).
- Ollinger, S. V., Aber, J. D., Reich, P. B. & Freuder, R. J. Interactive effects of nitrogen deposition, tropospheric ozone, elevated CO₂, and land use history on the carbon dynamics of northern hardwood forests. *Glob. Change Biol.* **8**, 545–562 (2002).
- Nicotra, A. B. et al. The evolution and functional significance of leaf shape in the angiosperms. *Funct. Plant Biol.* **38**, 535–552 (2011).
- Díaz, S. et al. The global spectrum of plant form and function. *Nature* **529**, 167–171 (2016).
- Baldocchi, D. D. et al. On the differential advantages of evergreenness and deciduousness in mediterranean oak woodlands: a flux perspective. *Ecol. Appl.* **20**, 1583–1597 (2010).
- Arora, V. K. & Boer, G. J. A parameterization of leaf phenology for the terrestrial ecosystem component of climate models. *Glob. Change Biol.* **11**, 39–59 (2005).
- Schweitzer, J. A. et al. Genetically based trait in a dominant tree affects ecosystem processes. *Ecol. Lett.* **7**, 127–134 (2004).
- Tian, F. et al. Coupling of ecosystem-scale plant water storage and leaf phenology observed by satellite. *Nat. Ecol. Evol.* **2**, 1428–1435 (2018).
- Méndez-Alonzo, R., Pineda-García, F., Paz, H., Rosell, J. A. & Olson, M. E. Leaf phenology is associated with soil water availability and xylem traits in a tropical dry forest. *Trees* **27**, 745–754 (2013).
- Givnish, T. J. Adaptive significance of evergreen vs. deciduous leaves: solving the triple paradox. *Silva Fenn.* **36**, 703–743 (2002).
- Axelrod, D. I. Origin of deciduous and evergreen habits in temperate forests. *Evolution* **20**, 1–15 (1966).
- Reich, P. B., Walters, M. B. & Ellsworth, D. S. From tropics to tundra: global convergence in plant functioning. *Proc. Natl Acad. Sci. USA* **94**, 13730–13734 (1997).
- Villar, R. & Merino, J. Comparison of leaf construction costs in woody species with differing leaf life-spans in contrasting ecosystems. *New Phytol.* **151**, 213–226 (2001).
- Chabot, B. F. & Hicks, D. J. The ecology of leaf life spans. *Annu. Rev. Ecol. Syst.* **13**, 229–259 (2003).
- Augusto, L. et al. Influences of evergreen gymnosperm and deciduous angiosperm tree species on the functioning of temperate and boreal forests. *Biol. Rev.* **90**, 444–466 (2015).
- Flo, V. et al. Climate and functional traits jointly mediate tree water-use strategies. *New Phytol.* **231**, 617–630 (2021).
- Lin, Y. S. et al. Optimal stomatal behaviour around the world. *Nat. Clim. Change* **5**, 459–464 (2015).
- Choat, B. et al. Global convergence in the vulnerability of forests to drought. *Nature* **491**, 752–755 (2012).
- Lusk, C. H., Wright, I. & Reich, P. B. Photosynthetic differences contribute to competitive advantage of evergreen angiosperm trees over evergreen conifers in productive habitats. *New Phytol.* **160**, 329–336 (2003).
- Mekonnen, Z. A., Riley, W. J., Randerson, J. T., Grant, R. F. & Rogers, B. M. Expansion of high-latitude deciduous forests driven by interactions between climate warming and fire. *Nat. Plants* **5**, 952–958 (2019).
- Baltzer, J. L. et al. Increasing fire and the decline of fire adapted black spruce in the boreal forest. *Proc. Natl Acad. Sci. USA* **118**, e2024872118 (2021).
- Mack, M. C. et al. Carbon loss from boreal forest wildfires offset by increased dominance of deciduous trees. *Science* **372**, 280–283 (2021).
- Kikuzawa, K. A cost-benefit analysis of leaf habit and leaf longevity of trees and their geographical pattern. *Am. Nat.* **138**, 1250–1263 (1991).
- Huechacona-Ruiz, A. H. et al. Mapping tree species deciduousness of tropical dry forests combining reflectance, spectral unmixing, and texture data from high-resolution imagery. *Forests* **11**, 1234 (2020).
- Sitch, S. et al. Evaluation of ecosystem dynamics, plant geography and terrestrial carbon cycling in the LPJ dynamic global vegetation model. *Glob. Change Biol.* **9**, 161–185 (2003).
- Woodward, F. I. & Williams, B. G. Climate and plant distribution at global and local scales. *Vegetatio* **69**, 189–197 (1987).
- Smith, B. et al. Implications of incorporating N cycling and N limitations on primary production in an individual-based dynamic vegetation model. *Biogeosciences* **11**, 2027–2054 (2014).
- Bondeau, A. et al. Modelling the role of agriculture for the 20th century global terrestrial carbon balance. *Glob. Change Biol.* **13**, 679–706 (2007).
- Gerten, D., Schaphoff, S., Haberlandt, U., Lucht, W. & Sitch, S. Terrestrial vegetation and water balance - hydrological evaluation of a dynamic global vegetation model. *J. Hydrol.* **286**, 249–270 (2004).
- Krinner, G. et al. A dynamic global vegetation model for studies of the coupled atmosphere-biosphere system. *Glob. Biogeochem. Cycles* **19**, GB1015 (2005).
- Sato, H., Itoh, A. & Kohyama, T. SEIB-DGVM: a new dynamic global vegetation model using a spatially explicit individual-based approach. *Ecol. Modell.* **200**, 279–307 (2007).
- Liang, J. et al. Positive biodiversity-productivity relationship predominant in global forests. *Science* **354**, aaf8957 (2016).
- Kattge, J. et al. TRY-a global database of plant traits. *Glob. Change Biol.* **17**, 2905–2935 (2011).
- Jucker, T. et al. Tallo: a global tree allometry and crown architecture database. *Glob. Change Biol.* <https://doi.org/10.1111/gcb.16302> (2022).
- Land Cover Classification Gridded Maps from 1992 to Present Derived From Satellite Observations* (Copernicus Climate Change Service (C3S) Climate Data Store (CDS), accessed 24 March 2023); <https://doi.org/10.24381/cds.006f2c9a>

42. Goldberg, D. E. The distribution of evergreen and deciduous trees relative to soil type: an example from the Sierra Madre, Mexico, and a general model. *Ecology* **63**, 942–951 (1982).
43. Reichert, J. M. et al. Estimation of water retention and availability in soils of Rio Grande do Sul. *Rev. Bras. Cienc. Solo* **33**, 1547–1560 (2009).
44. Duong, T. T. T., Penfold, C. & Marschner, P. Amending soils of different texture with six compost types: impact on soil nutrient availability, plant growth and nutrient uptake. *Plant Soil* **354**, 197–209 (2012).
45. Yang, J. et al. Leaf form-climate relationships on the global stage: an ensemble of characters. *Glob. Ecol. Biogeogr.* **24**, 1113–1125 (2015).
46. Allen, C. D. et al. A global overview of drought and heat-induced tree mortality reveals emerging climate change risks for forests. *Ecol. Manage.* **259**, 660–684 (2010).
47. Crowther, T. W. et al. Mapping tree density at a global scale. *Nature* **525**, 201–205 (2015).
48. Gatti, R. C. et al. The number of tree species on Earth. *Proc. Natl Acad. Sci. USA* **119**, (2022).
49. Santoro, M. & Cartus, O. *ESA Biomass Climate Change Initiative (Biomass_cci): Global Datasets of Forest Above-ground Biomass for the Years 2010, 2017 and 2018 v.3* (NERC EDS Centre for Environmental Data Analysis, 2021).
50. Karger, D. N. et al. Climatologies at high resolution for the earth's land surface areas. *Sci. Data* **4**, 170122 (2017).
51. Reich, P. B. et al. Even modest climate change may lead to major transitions in boreal forests. *Nature* **608**, 540–545 (2022).
52. Elsen, P. R. et al. Accelerated shifts in terrestrial life zones under rapid climate change. *Glob. Change Biol.* **28**, 918–935 (2022).
53. Graham, R. L., Turner, M. G. & Dale, V. H. How increasing CO₂ and climate change affect forests. *Bioscience* **40**, 575–587 (1990).
54. Keenan, T., Maria Serra, J., Lloret, F., Ninyerola, M. & Sabate, S. Predicting the future of forests in the Mediterranean under climate change, with niche- and process-based models: CO₂ matters! *Glob. Change Biol.* **17**, 565–579 (2011).
55. Liang, J. et al. Co-limitation towards lower latitudes shapes global forest diversity gradients. *Nat. Ecol. Evol.* **6**, 1423–1437 (2022).
56. Batjes, N. H., Ribeiro, E. & Van Oostrum, A. Standardised soil profile data to support global mapping and modelling (WoSIS snapshot 2019). *Earth Syst. Sci. Data* **12**, 299–320 (2020).
57. Condit, R., Engelbrecht, B. M. J., Pino, D., Pérez, R. & Turner, B. L. Species distributions in response to individual soil nutrients and seasonal drought across a community of tropical trees. *Proc. Natl Acad. Sci. USA* **110**, 5064–5068 (2013).
58. Álvarez-Yépiz, J. C. et al. Resource partitioning by evergreen and deciduous species in a tropical dry forest. *Oecologia* **183**, 607–618 (2017).
59. Aerts, R. The advantages of being evergreen. *Trends Ecol. Evol.* **10**, 402–407 (1995).
60. Ouédraogo, D.-Y. et al. The determinants of tropical forest deciduousness: disentangling the effects of rainfall and geology in central Africa. *J. Ecol.* **104**, 924–935 (2016).
61. Chave, J. et al. Towards a worldwide wood economics spectrum. *Ecol. Lett.* **12**, 351–366 (2009).
62. Simard, M., Pinto, N., Fisher, J. B. & Baccini, A. Mapping forest canopy height globally with spaceborne lidar. *J. Geophys. Res. Biogeosci.* **116**, G04021 (2011).
63. Sitch, S. et al. Recent trends and drivers of regional sources and sinks of carbon dioxide. *Biogeosciences* **12**, 653–679 (2015).
64. Pugh, T. A. M. et al. Understanding the uncertainty in global forest carbon turnover. *Biogeosciences* **17**, 3961–3989 (2020).
65. Hansen, M. C. et al. High-resolution global maps of 21st-century forest cover change. *Science* **342**, 850–853 (2013).
66. Cayuela, L., Granzow-de la Cerda, Í., Albuquerque, F. S. & Golicher, D. J. Taxonstand: an R package for species names standardisation in vegetation databases. *Methods Ecol. Evol.* **3**, 1078–1083 (2012).
67. Hengl, T. et al. SoilGrids250m: global gridded soil information based on machine learning. *PLoS ONE* **12**, e0169748 (2017).
68. Gorelick, N. et al. Google Earth Engine: planetary-scale geospatial analysis for everyone. *Remote Sens. Environ.* **202**, 18–27 (2017).
69. Strobl, C., Boulesteix, A. L., Kneib, T., Augustin, T. & Zeileis, A. Conditional variable importance for random forests. *BMC Bioinformatics* **9**, 307 (2008).
70. Simin, C., Rongqun, Z., Wenling, C. & Hui, Y. Band selection of hyperspectral images based on Bhattacharyya distance. *WSEAS Trans. Inf. Sci. Appl.* **6**, 1165–1175 (2009).
71. Ning, J., Zhang, L., Zhang, D. & Wu, C. Interactive image segmentation by maximal similarity based region merging. *Pattern Recognit.* **43**, 445–456 (2010).
72. Choi, E. & Lee, C. Feature extraction based on the Bhattacharyya distance. *Pattern Recognit.* **36**, 1703–1709 (2003).
73. El Merabet, Y. et al. Maximal similarity based region classification method through local image region descriptors and Bhattacharyya coefficient-based distance: application to horizon line detection using wide-angle camera. *Neurocomputing* **265**, 28–41 (2017).
74. Li, J. Assessing the accuracy of predictive models for numerical data: not r nor r^2 , why not? Then what? *PLoS ONE* **12**, e0183250 (2017).
75. Bhattacharyya, A. On a measure of divergence between two multinomial populations. *Sankhyā Indian J. Stat.* **7**, 401–406 (1946).
76. Ma, H. et al. The global distribution and environmental drivers of aboveground versus belowground plant biomass. *Nat. Ecol. Evol.* **5**, 1110–1122 (2021).
77. Breiman, L., Friedman, J., Stone, C. J. & Olshen, R. A. *Classification Algorithms and Regression Trees* (Chapman & Hall, 1984).
78. Hijmans, R. J. et al. Package geosphere (CRAN, 2019).
79. van den Hoogen, J. et al. Soil nematode abundance and functional group composition at a global scale. *Nature* **572**, 194–198 (2019).
80. Olson, D. M. et al. Terrestrial ecoregions of the world: a new map of life on Earth. *Bioscience* **51**, 933–938 (2001).
81. Tuanmu, M. N. & Jetz, W. A global 1-km consensus land-cover product for biodiversity and ecosystem modelling. *Glob. Ecol. Biogeogr.* **23**, 1031–1045 (2014).
82. Heiberger, R. M. *HH: Statistical Analysis and Data Display: Heiberger and Holland* (CRAN, 2020).
83. Erin, L. et al. h2o: R Interface for the 'H2O' Scalable Machine Learning Platform. R package v.3.32.0.2 (GitHub, 2020).

Acknowledgements

We thank the Global Forest Biodiversity Initiative (GFBi) for establishing the data standards and collaborative framework; TRY database managers and countless researchers who contributed open-access data. This work was supported by grants to C.M.Z. from the Ambizione fellowship (PZ00P3_193646), L.M. from the China Scholarship Council, and T.W.C. from DOB Ecology and the Bernina Foundation. O.B. acknowledges funding from the Romania National Council for Higher Education, CNFIS, project number CNFIS-FDI-2023-F-0579. We thank the French National Forest Inventory (NFI campaigns; raw data 2005 and subsequent annual surveys were downloaded by GFBi at <https://inventaire-forestier.ign.fr/spip.php?rubrique159>, accessed on 1 January 2015) and the Italian Forest Inventory (NFI campaigns; raw data 2005 and subsequent surveys were downloaded by GFBi at <https://inventarioforestale.org/>, accessed on 27 April 2019); G.A. was supported by the Italian National Recovery Plan through the National Biodiversity Future Center. M.B. thanks the BOS Foundation,

the Indonesian Institute of Sciences (LIPI), the Direktorat Fasilitas Organisasi Politik dan Masyarakat, Departemen Dalam Negeri, and the BKSDA Palangkaraya for permission to work in the MAWAS region of Indonesia, and acknowledges funding from The American Society of Primatologists, the Duke University Graduate School, the L.S.B. Leakey Foundation, the National Science Foundation (Grant No. 0452995) and the Wenner-Gren Foundation for Anthropological Research (Grant No. 7330). K.S. thanks the IBL for supporting this work by internal funds under project no. 261509 "AFTER FBS - maintenance of ForBioSensing project performance indicators". Plot data collection from Białowieża Forest was supported by Project LIFE+ ForBioSensing (contract number LIFE13ENV/PL/000048) and Poland's National Fund for Environmental Protection and Water Management (contract number 485/2014/WN10/OPNMLF/D). Plot data collection from Central Siberia was supported by Russian Science Foundation project no 21-46-07002. Contributions from ForestPlots.net and RAINFOR were supported by numerous sources, including the European Research Council (ERC Advanced Grant 291585 - 'T-FORCES'), the Gordon and Betty Moore Foundation (1656 'RAINFOR'), the Natural Environment Research Council (including NE/B503384/1, NE/NO12542/1) and the Royal Society (ICA/R1/180100 - 'FORAMA'). We also thank the National Council for Science and Technology Development of Brazil (CNPq) for support to the Cerrado/Amazonia Transition Long-Term Ecology Project (PELD/441244/2016-5). J.D. was funded by the Czech Science Foundation (project no. 21-26883S). We are grateful to all the ministries and agencies from the Government of Spain and to all the people that supported the collection, compilation, and coordination of forest inventory data, also including the Spanish National Forest Inventories. Sergio de Miguel benefitted from a Serra-Hunter fellowship provided by the Government of Catalonia (Generalitat de Catalunya). J.C.S. considers this work a contribution to Center for Ecological Dynamics in a Novel Biosphere (ECONOVO), funded by the Danish National Research Foundation (grant DNR173) and his VILLUM Investigator project 'Biodiversity Dynamics in a Changing World', funded by VILLUM FONDEN (grant 16549). The exploratory plots of FunDivEUROPE were established through funding from the European Union Seventh Framework Programme FP7/2007-2013 under Grant 265171. Plot data collection from the Brazilian Atlantic Forest was supported by the State of São Paulo Research Foundation/FAPESP (BIOTA Grants 2003/12595-7; 2012/51509-8 and 2012/51872-5) and by the Brazilian National Research Council/CNPq (PELD Grant 403710/2012-0). H.V. was funded by FCT - Portuguese Foundation for Science and Technology, project UIDB/04033/2020 and ICNF-Instituto da Conservação da Natureza, Portugal, National Forest Inventory. Financial support for the Monafor network in Mexico came from many projects, including the National Forestry Commission (CONAFOR), Council of Science and Technology of the State of Durango (COCYTED), the Natural Environment Research Council, UK (NERC; NE/T011084/1), and local support of Ejidos and Comunidades. We also directly acknowledge the use of data drawn from the Natural Forest plot data collected between January 2009 and

March 2014 by the LUCAS programme for the New Zealand Ministry for the Environment. Data were sourced via the NZ National Vegetation Survey (NVS) Databank.

Author contributions

H.M. and C.M.Z. conceived, developed and wrote the paper, with input from T.W.C., L.M., D.S.M., S.S.R., and Y.Z. H.M. performed the analyses with assistance from C.M.Z., Y.Z., J.v.d.H., D.S.M. and L.M. All authors contributed to data collection and/or curation and reviewed and/or provided input on the manuscript.

Funding

Open access funding provided by Swiss Federal Institute of Technology Zurich.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41477-023-01543-5>.

Correspondence and requests for materials should be addressed to Haozhi Ma.

Peer review information *Nature Plants* thanks Helena Vallicrosa and the other, anonymous, reviewer(s) for their contribution to the peer review of this work.

Reprints and permissions information is available at www.nature.com/reprints.

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2023

Haozhi Ma¹✉, Thomas W. Crowther¹, Lidong Mo¹, Daniel S. Maynard^{1,209}, Susanne S. Renner¹², Johan van den Hoogen¹, Yibiao Zou¹, Jingjing Liang³, Sergio de-Miguel^{4,5}, Gert-Jan Nabuurs⁶, Peter B. Reich^{7,8,9}, Ülo Niinemets¹⁰, Meinrad Abegg¹¹, Yves C. Adou Yao¹², Giorgio Alberti^{13,14}, Angelica M. Almeyda Zambrano¹⁵, Braulio Vilchez Alvarado¹⁶, Esteban Alvarez-Dávila¹⁷, Patricia Alvarez-Loayza¹⁸, Luciana F. Alves¹⁹, Christian Ammer²⁰, Clara Antón-Fernández²¹, Alejandro Araujo-Murakami²², Luzmila Arroyo²², Valerio Avitabile²³, Gerardo A. Aymard^{24,25}, Timothy R. Baker²⁶, Radomir Batázy²⁷, Olaf Banki²⁸, Jorcely G. Barroso²⁹, Meredith L. Bastian^{30,31}, Jean-Francois Bastin³², Luca Birigazzi³³, Philippe Birnbaum^{34,35,36}, Robert Bitariho³⁷, Pascal Boeckx³⁸, Frans Bongers³⁹, Olivier Bouriaud³⁹, Pedro H. S. Brancalion⁴⁰, Susanne Brandt⁴¹, Francis Q. Brearley⁴², Roel Brienens²⁶, Eben N. Broadbent¹⁵, Helge Bruehlheide^{43,44}, Filippo Bussotti⁴⁵, Roberto Cazzolla Gatti⁴⁶,

Ricardo G. César⁴⁰, Goran Cesljar⁴⁷, Robin Chazdon^{48,49}, Han Y. H. Chen⁵⁰, Chelsea Chisholm¹, Hyunkook Cho⁵¹, Emil Cienciala^{52,53}, Connie Clark⁵⁴, David Clark⁵⁵, Gabriel D. Colletta⁵⁶, David A. Coomes⁵⁷, Fernando Cornejo Valverde⁵⁸, José J. Corral-Rivas⁵⁹, Philip M. Crim^{60,61}, Jonathan R. Cumming⁶⁰, Selvadurai Dayanandan⁶², André L. de Gasper⁶³, Mathieu Decuyper⁶, Géraldine Derroire⁶⁴, Ben DeVries⁶⁵, Ilija Djordjevic⁶⁶, Jiri Dolezal^{67,68}, Aurélie Dourdain⁶⁴, Nestor Laurier Engone Obiang⁶⁹, Brian J. Enquist^{70,71}, Teresa J. Eyre⁷², Adandé Belarmain Fandohan⁷³, Tom M. Fayle^{74,75}, Ted R. Feldpausch⁷⁶, Leandro V. Ferreira⁷⁷, Leena Finér⁷⁸, Markus Fischer⁷⁹, Christine Fletcher⁸⁰, Jonas Fridman⁸¹, Lorenzo Frizzera⁸², Javier G. P. Gamarra⁸³, Damiano Gianelle⁸², Henry B. Glick⁸⁴, David J. Harris⁸⁵, Andrew Hector⁸⁶, Andreas Hemp⁸⁷, Geerten Hengeveld⁶, Bruno Hérault^{88,89}, John L. Herbohn⁹⁰, Martin Herold⁹¹, Annika Hillers^{92,93}, Eurídice N. Honorio Coronado⁹⁴, Cang Hui^{95,96}, Thomas T. Ibanez⁹⁷, Iêda Amaral⁹⁸, Nobuo Imai⁹⁹, Andrzej M. Jagodziński^{100,101}, Bogdan Jaroszewicz¹⁰², Vivian Kvist Johannsen¹⁰³, Carlos A. Joly¹⁰⁴, Tommaso Jucker¹⁰⁵, Ilbin Jung⁵¹, Viktor Karminov¹⁰⁶, Kuswata Kartawinata¹⁸, Elizabeth Kearsley¹⁰⁷, David Kenfack¹⁰⁸, Deborah K. Kennard¹⁰⁹, Sebastian Kepfer-Rojas¹⁰³, Gunnar Keppel¹¹⁰, Mohammed Latif Khan¹¹¹, Timothy J. Killeen²², Hyun Seok Kim^{112,113,114,115}, Kanehiro Kitayama¹¹⁶, Michael Köhl¹¹⁷, Henn Korjus¹¹⁸, Florian Kraxner¹¹⁹, Dmitry Kucher¹²⁰, Diana Laarmann¹¹⁸, Mait Lang¹¹⁸, Simon L. Lewis^{26,121}, Huicui Lu¹²², Natalia V. Lukina¹²³, Brian S. Maitner⁷⁰, Yadvinder Malhi¹²⁴, Eric Marcon¹²⁵, Beatriz Schwantes Marimon¹²⁶, Ben Hur Marimon-Junior¹²⁶, Andrew R. Marshall^{90,127,128}, Emanuel H. Martin¹²⁹, Jorge A. Meave¹³⁰, Omar Melo-Cruz¹³¹, Casimiro Mendoza¹³², Cory Merow⁴⁸, Abel Monteagudo Mendoza^{133,134}, Vanessa S. Moreno⁴⁰, Sharif A. Mukul^{90,135}, Philip Mundhenk¹¹⁷, María Guadalupe Nava-Miranda^{136,137}, David Neill¹³⁸, Victor J. Neldner⁷², Radovan V. Nevenic⁶⁶, Michael R. Ngugi⁷², Pascal A. Niklaus¹³⁹, Jacek Oleksyn¹⁰⁰, Petr Ontikov¹⁰⁶, Edgar Ortiz-Malavasi¹⁶, Yude Pan¹⁴⁰, Alain Paquette¹⁴¹, Alexander Parada-Gutierrez²², Elena I. Parfenova¹⁴², Minjee Park^{3,112}, Marc Parren¹⁴³, Narayanaswamy Parthasarathy¹⁴⁴, Pablo L. Peri¹⁴⁵, Sebastian Pfautsch¹⁴⁶, Oliver L. Phillips²⁶, Nicolas Picard¹⁴⁷, Maria Teresa F. Piedade¹⁴⁸, Daniel Piotto¹⁴⁹, Nigel C. A. Pitman¹⁵⁰, Irina Mendoza-Polo¹⁵¹, Axel D. Poulsen⁸⁵, John R. Poulsen^{54,152}, Hans Pretzsch^{153,154}, Freddy Ramirez Arevalo¹⁵⁵, Zorayda Restrepo-Correa¹⁵⁶, Mirco Rodeghiero^{82,157}, Samir G. Rolim¹⁴⁹, Anand Roopsind¹⁵⁸, Francesco Rovero^{159,160}, Ervan Rutishauser¹⁶¹, Purabi Saikia¹⁶², Christian Salas-Eljatib^{163,164,165}, Philippe Saner¹⁶⁶, Peter Schall²⁰, Mart-Jan Schelhaas⁶, Dmitry Schepaschenko^{119,167}, Michael Scherer-Lorenzen¹⁶⁸, Bernhard Schmid¹⁶⁹, Jochen Schöngart¹⁴⁸, Eric B. Searle¹⁴¹, Vladimír Seben¹⁷⁰, Josep M. Serra-Diaz^{171,172}, Douglas Sheil^{143,173}, Anatoly Z. Shvidenko¹¹⁹, Javier E. Silva-Espejo¹⁷⁴, Marcos Silveira¹⁷⁵, James Singh¹⁷⁶, Plinio Sist⁸⁸, Ferry Slik¹⁷⁷, Bonaventure Sonké¹⁷⁸, Alexandre F. Souza¹⁷⁹, Stanislaw Miścicki¹⁸⁰, Krzysztof J. Stereńczak²⁷, Jens-Christian Svenning^{181,182}, Miroslav Svoboda¹⁸³, Ben Swanepoel¹⁸⁴, Natalia Targhetta¹⁴⁸, Nadja Tchebakova¹⁴², Hans ter Steege^{28,185}, Raquel Thomas¹⁸⁶, Elena Tikhonova¹²³, Peter M. Umunay¹⁸⁷, Vladimir A. Usoltsev¹⁸⁸, Renato Valencia¹⁸⁹, Fernando Valladares¹⁹⁰, Fons van der Plas¹⁹¹, Tran Van Do¹⁹², Michael E. van Nuland¹⁹³, Rodolfo M. Vasquez¹³³, Hans Verbeek¹⁰⁷, Helder Viana^{194,195}, Alexander C. Vibrans^{63,196}, Simone Vieira¹⁹⁷, Klaus von Gadow¹⁹⁸, Hua-Feng Wang¹⁹⁹, James V. Watson²⁰⁰, Gijsbert D. A. Werner²⁰¹, Bertil Westerlund⁸¹, Susan K. Wiser²⁰², Florian Wittmann²⁰³, Hannsjoerg Woell²⁰⁴, Verginia Wortel²⁰⁵, Roderick Zagt²⁰⁶, Tomasz Zawila-Niedzwiecki²⁰⁷, Chunyu Zhang²⁰⁸, Xiuhai Zhao²⁰⁸, Mo Zhou³, Zhi-Xin Zhu¹⁹⁹, Irie C. Zo-Bi⁸⁹ & Constantin M. Zohner¹

¹Institute of Integrative Biology, ETH Zurich (Swiss Federal Institute of Technology), Zurich, Switzerland. ²Department of Biology, Washington University, Saint Louis, MO, USA. ³Department of Forestry and Natural Resources, Purdue University, West Lafayette, IN, USA. ⁴Department of Agricultural and Forest Sciences and Engineering, University of Lleida, Lleida, Spain. ⁵Joint Research Unit CTFC - AGROTECNIO - CERCA, Solsona, Spain. ⁶Wageningen University and Research, Wageningen, the Netherlands. ⁷Department of Forest Resources, University of Minnesota, St. Paul, MN, USA. ⁸Hawkesbury Institute for the Environment, Western Sydney University, Penrith, New South Wales, Australia. ⁹Institute for Global Change Biology, and School for Environment and Sustainability, University of Michigan, Ann Arbor, MI, USA. ¹⁰Chair of Crop Science and Plant Biology, Estonian University of Life Sciences, Tartu, Estonia. ¹¹Swiss Federal Institute for Forest, Snow and Landscape Research WSL, Birmensdorf, Switzerland. ¹²UFR Biosciences, University Félix Houphouët-Boigny, Abidjan, Côte d'Ivoire. ¹³Department of Agricultural, Food, Environmental and Animal Sciences, University of Udine, Udine, Italy. ¹⁴National Biodiversity Future Center (NBFC), Palermo, Italy. ¹⁵Spatial Ecology and Conservation Laboratory, Department of Tourism, Recreation and Sport Management, University of Florida, Gainesville, FL, USA. ¹⁶Forestry School, Tecnológico de Costa Rica TEC, Cartago, Costa Rica. ¹⁷Fundacion ConVida, Universidad Nacional Abierta y a Distancia, UNAD, Medellín, Colombia. ¹⁸Negaunee Integrative Research Center, Field Museum of Natural History, Chicago, IL, USA. ¹⁹Center for Tropical Research, Institute of the Environment and Sustainability, UCLA, Los Angeles, CA, USA. ²⁰Silviculture and Forest Ecology of the Temperate Zones, University of Göttingen, Göttingen, Germany. ²¹Division of Forest and Forest Resources, Norwegian Institute of Bioeconomy Research (NIBIO), Ås, Norway. ²²Museo de Historia natural Noel Kempff Mercado, Santa Cruz, Bolivia. ²³European Commission, Joint Research Center, Ispra, Italy. ²⁴UNELLEZ-Guanare, Programa de Ciencias del Agro y el Mar, Herbario Universitario (PORT), Portuguesa, Venezuela. ²⁵Compensation International S. A. Ci Progress-GreenLife, Bogotá, D.C., Colombia. ²⁶School of Geography, University of Leeds, Leeds, UK. ²⁷Department of Geomatics, Forest Research Institute, Raszyn, Poland. ²⁸Naturalis Biodiversity Center, Leiden, the Netherlands. ²⁹Centro Multidisciplinar, Universidade Federal do Acre, Rio Branco, Brazil. ³⁰Proceedings of the National Academy of Sciences, Washington, DC, USA. ³¹Department of Evolutionary Anthropology, Duke University, Durham, NC, USA. ³²TERRA Teach and Research Centre, Gembloux Agro Bio-Tech, University of Liege, Liege, Belgium. ³³Forestry Consultant, Grosseto, Italy.

³⁴Institut Agronomique néo-Calédonien (IAC), Nouméa, New Caledonia. ³⁵AMAP, Univ. Montpellier, Montpellier, France. ³⁶CIRAD, CNRS, INRAE, IRD, Montpellier, France. ³⁷Institute of Tropical Forest Conservation, Mbarara University of Sciences and Technology, Mbarara, Uganda. ³⁸Isotope Bioscience Laboratory - ISOFYS, Ghent University, Ghent, Belgium. ³⁹Ștefan cel Mare University of Suceava, Suceava, Romania. ⁴⁰Department of Forest Sciences, Luiz de Queiroz College of Agriculture, University of São Paulo, Piracicaba, Brazil. ⁴¹Bavarian State Institute of Forestry, Freising, Germany. ⁴²Department of Natural Sciences, Manchester Metropolitan University, Manchester, UK. ⁴³Institute of Biology, Geobotany and Botanical Garden, Martin Luther University Halle-Wittenberg, Halle-Wittenberg, Germany. ⁴⁴German Centre for Integrative Biodiversity Research (iDiv) Halle-Jena-Leipzig, Leipzig, Germany. ⁴⁵Department of Agriculture, Food, Environment and Forest (DAGRI), University of Firenze, Florence, Italy. ⁴⁶Department of Biological, Geological, and Environmental Sciences, University of Bologna, Bologna, Italy. ⁴⁷Department of Spatial Regulation GIS and Forest Policy, Institute of Forestry, Belgrade, Serbia. ⁴⁸Department of Ecology and Evolutionary Biology, University of Connecticut, Storrs, CT, USA. ⁴⁹Tropical Forest and People Research Centre, University of the Sunshine Coast, Sippy Downs, Queensland, Australia. ⁵⁰Faculty of Natural Resources Management, Lakehead University, Thunder Bay, Ontario, Canada. ⁵¹Division of Forest Resources Information, Korea Forest Promotion Institute, Seoul, South Korea. ⁵²IFER - Institute of Forest Ecosystem Research, Jilove u Prahy, Czech Republic. ⁵³Global Change Research Institute CAS, Brno, Czech Republic. ⁵⁴Nicholas School of the Environment, Duke University, Durham, NC, USA. ⁵⁵Department of Biology, University of Missouri-St Louis, St. Louis, MO, USA. ⁵⁶Programa de Pós-graduação em Biologia Vegetal, Instituto de Biologia, Universidade Estadual de Campinas, Campinas, Brazil. ⁵⁷Department of Plant Sciences and Conservation Research Institute, University of Cambridge, Cambridge, UK. ⁵⁸Andes to Amazon Biodiversity Program, Madre de Dios, Peru. ⁵⁹Facultad de Ciencias Forestales y Ambientales, Universidad Juárez del Estado de Durango, Durango, Mexico. ⁶⁰Department of Biology, West Virginia University, Morgantown, WV, USA. ⁶¹Department of Physical and Biological Sciences, The College of Saint Rose, Albany, NY, USA. ⁶²Biology Department, Centre for Structural and Functional Genomics, Concordia University, Montreal, Quebec, Canada. ⁶³Natural Science Department, Universidade Regional de Blumenau, Blumenau, Brazil. ⁶⁴Cirad, UMR EcoFoG (AgroParisTech, CNRS, INRAE, Université des Antilles Université de la Guyane), Campus Agronomique, Kourou, French Guiana. ⁶⁵Department of Geography, Environment and Geomatics, University of Guelph, Guelph, Ontario, Canada. ⁶⁶Institute of Forestry, Belgrade, Serbia. ⁶⁷Institute of Botany, The Czech Academy of Sciences, Třeboň, Czech Republic. ⁶⁸Department of Botany, Faculty of Science, University of South Bohemia, České Budějovice, Czech Republic. ⁶⁹IRET, Herbarier National du Gabon (CENAREST), Libreville, Gabon. ⁷⁰Department of Ecology and Evolutionary Biology, University of Arizona, Tucson, AZ, USA. ⁷¹The Santa Fe Institute, Santa Fe, NM, USA. ⁷²Queensland Herbarium and Biodiversity Science, Department of Environment and Science, Toowong, Queensland, Australia. ⁷³Ecole de Foresterie et Ingénierie du Bois, Université Nationale d'Agriculture, Kétou, Benin. ⁷⁴School of Biological and Behavioural Sciences, Queen Mary University of London, London, UK. ⁷⁵Biology Centre of the Czech Academy of Sciences, Institute of Entomology, Ceske Budejovice, Czech Republic. ⁷⁶Geography, College of Life and Environmental Sciences, University of Exeter, Exeter, UK. ⁷⁷Museu Paraense Emílio Goeldi. Coordenação de Ciências da Terra e Ecologia, Belém, Pará, Brasil. ⁷⁸Natural Resources Institute Finland (Luke), Joensuu, Finland. ⁷⁹Institute of Plant Sciences, University of Bern, Bern, Switzerland. ⁸⁰Forest Research Institute Malaysia, Kuala Lumpur, Malaysia. ⁸¹Department of Forest Resource Management, Swedish University of Agricultural Sciences SLU, Umea, Sweden. ⁸²Research and Innovation Center, Fondazione Edmund Mach, San Michele All'adige, Italy. ⁸³Forestry Division, Food and Agriculture Organization of the United Nations, Rome, Italy. ⁸⁴Glick Designs LLC, Hadley, MA, USA. ⁸⁵Royal Botanic Garden Edinburgh, Edinburgh, UK. ⁸⁶Department of Biology, University of Oxford, Oxford, UK. ⁸⁷Department of Plant Systematics, University of Bayreuth, Bayreuth, Germany. ⁸⁸Cirad, UPR Forêts et Sociétés, University of Montpellier, Montpellier, France. ⁸⁹Department of Forestry and Environment, National Polytechnic Institute (INP-HB), Yamoussoukro, Côte d'Ivoire. ⁹⁰Forest Research Institute, University of the Sunshine Coast, Sippy Downs, Queensland, Australia. ⁹¹Helmholtz GFZ German Research Centre for Geosciences, Remote Sensing and Geoinformatics Section, Telegrafenberg, Potsdam, Germany. ⁹²Centre for Conservation Science, The Royal Society for the Protection of Birds, Sandy, UK. ⁹³Wild Chimpanzee Foundation, Liberia Office, Monrovia, Liberia. ⁹⁴Instituto de Investigaciones de la Amazonia Peruana, Iquitos, Peru. ⁹⁵Centre for Invasion Biology, Department of Mathematical Sciences, Stellenbosch University, Stellenbosch, South Africa. ⁹⁶Theoretical Ecology Unit, African Institute for Mathematical Sciences, Cape Town, South Africa. ⁹⁷AMAP, Univ Montpellier, CIRAD, CNRS, INRAE, IRD, Montpellier, France. ⁹⁸National Institute of Amazonian Research, Manaus, Brazil. ⁹⁹Department of Forest Science, Tokyo University of Agriculture, Tokyo, Japan. ¹⁰⁰Institute of Dendrology, Polish Academy of Sciences, Kórnik, Poland. ¹⁰¹Department of Game Management and Forest Protection, Poznań University of Life Sciences, Poznań, Poland. ¹⁰²Faculty of Biology, Białowieża Geobotanical Station, University of Warsaw, Białowieża, Poland. ¹⁰³Department of Geosciences and Natural Resource Management, University of Copenhagen, Copenhagen, Denmark. ¹⁰⁴Department of Plant Biology, Institute of Biology, University of Campinas, UNICAMP, Campinas, Brazil. ¹⁰⁵School of Biological Sciences, University of Bristol, Bristol, UK. ¹⁰⁶Forestry Faculty, Mytischki Branch of Bauman Moscow State Technical University, Mytischki, Russian Federation. ¹⁰⁷CAVELab-Computational and Applied Vegetation Ecology, Department of Environment, Ghent University, Ghent, Belgium. ¹⁰⁸CTFS-ForestGEO, Smithsonian Tropical Research Institute, Balboa, Panama. ¹⁰⁹Department of Physical and Environmental Sciences, Colorado Mesa University, Grand Junction, CO, USA. ¹¹⁰UniSA STEM and Future Industries Institute, University of South Australia, Adelaide, South Australia, Australia. ¹¹¹Department of Botany, Dr Harisingh Gour Vishwavidyalaya (A Central University), Sagar, India. ¹¹²Department of Agriculture, Forestry and Bioresources, Seoul National University, Seoul, South Korea. ¹¹³Interdisciplinary Program in Agricultural and Forest Meteorology, Seoul National University, Seoul, South Korea. ¹¹⁴National Center for Agro Meteorology, Seoul, South Korea. ¹¹⁵Research Institute for Agriculture and Life Sciences, Seoul National University, Seoul, South Korea. ¹¹⁶Graduate School of Agriculture, Kyoto University, Kyoto, Japan. ¹¹⁷Institute for World Forestry, University of Hamburg, Hamburg, Germany. ¹¹⁸Institute of Forestry and Engineering, Estonian University of Life Sciences, Tartu, Estonia. ¹¹⁹Biodiversity and Natural Resources Program, International Institute for Applied Systems Analysis, Laxenburg, Austria. ¹²⁰Peoples Friendship University of Russia (RUDN University), Moscow, Russian Federation. ¹²¹Department of Geography, University College London, London, UK. ¹²²Faculty of Forestry, Qingdao Agricultural University, Qingdao, China. ¹²³Center for Forest Ecology and Productivity, Russian Academy of Sciences, Moscow, Russian Federation. ¹²⁴Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, UK. ¹²⁵AgroParisTech, UMR-AMAP, Cirad, CNRS, INRAE, IRD, Université de Montpellier, Montpellier, France. ¹²⁶Departamento de Ciências Biológicas, Universidade do Estado de Mato Grosso, Nova Xavantina, Brazil. ¹²⁷Department of Environment and Geography, University of York, York, UK. ¹²⁸Flamingo Land Ltd, Kirby Misperton, UK. ¹²⁹Department of Wildlife Management, College of African Wildlife Management, Mweka, Tanzania. ¹³⁰Departamento de Ecología y Recursos Naturales, Facultad de Ciencias, Universidad Nacional Autónoma de México, Mexico City, Mexico. ¹³¹Universidad del Tolima, Ibagué, Colombia. ¹³²Colegio de Profesionales Forestales de Cochabamba, Cochabamba, Bolivia. ¹³³Jardín Botánico de Missouri, Pasco, Peru. ¹³⁴Universidad Nacional de San Antonio Abad del Cusco, Cusco, Peru. ¹³⁵Department of Environment and Development Studies, United International University, Dhaka, Bangladesh. ¹³⁶Instituto de Silvicultura e Industria de la Madera, Universidad Juárez del Estado de Durango, Durango, Mexico. ¹³⁷Programa de doctorado en Ingeniería para el desarrollo rural y civil, Escuela de Doctorado Internacional de la Universidad de Santiago de Compostela (EDIUS), Santiago de Compostela, Spain. ¹³⁸Universidad Estatal Amazónica, Puyo, Pastaza, Ecuador. ¹³⁹Department of Evolutionary Biology and Environmental Studies, University of Zürich, Zurich, Switzerland. ¹⁴⁰Climate, Fire, and Carbon Cycle Sciences, USDA Forest Service, Durham, NC, USA. ¹⁴¹Centre for Forest Research, Université du Québec à Montréal, Montréal, Québec, Canada.

¹⁴²V. N. Sukachev Institute of Forest, FRC KSC, Siberian Branch of the Russian Academy of Sciences, Krasnoyarsk, Russian Federation. ¹⁴³Forest Ecology and Forest Management Group, Wageningen University and Research, Wageningen, the Netherlands. ¹⁴⁴Department of Ecology and Environmental Sciences, Pondicherry University, Puducherry, India. ¹⁴⁵Instituto Nacional de Tecnología Agropecuaria (INTA), Universidad Nacional de la Patagonia Austral (UNPA), Consejo Nacional de Investigaciones Científicas y Técnicas (CONICET), Río Gallegos, Argentina. ¹⁴⁶School of Social Sciences (Urban Studies), Western Sydney University, Penrith, New South Wales, Australia. ¹⁴⁷GIP Ecofor, Paris, France. ¹⁴⁸Instituto Nacional de Pesquisas da Amazônia, Manaus, Brazil. ¹⁴⁹Laboratório de Dendrologia e Silvicultura Tropical, Centro de Formação em Ciências Agroflorestais, Universidade Federal do Sul da Bahia, Itabuna, Brazil. ¹⁵⁰Field Museum of Natural History, Chicago, IL, USA. ¹⁵¹Jardín Botánico de Medellín, Medellín, Colombia. ¹⁵²The Nature Conservancy, Boulder, CO, USA. ¹⁵³Chair for Forest Growth and Yield Science, Department of Life Science Systems, TUM School for Life Sciences, Technical University of Munich, Freising, Germany. ¹⁵⁴Sustainable Forest Management Research Institute iuFOR, University Valladolid, Valladolid, Spain. ¹⁵⁵Universidad Nacional de la Amazonía Peruana, Iquitos, Peru. ¹⁵⁶Servicios Ecosistémicos y Cambio Climático (SECC), Fundación Con Vida and Corporación COL-TREE, Medellín, Colombia. ¹⁵⁷Centro Agricoltura, Alimenti, Ambiente, University of Trento, San Michele All'adige, Italy. ¹⁵⁸Center for Natural Climate Solutions, Conservation International, Arlington, VA, USA. ¹⁵⁹Department of Biology, University of Florence, Florence, Italy. ¹⁶⁰Tropical Biodiversity, MUSE - Museo delle Scienze, Trento, Italy. ¹⁶¹Info Flora, Geneva, Switzerland. ¹⁶²Department of Environmental Sciences, Central University of Jharkhand, Ranchi, Jharkhand, India. ¹⁶³Centro de Modelación y Monitoreo de Ecosistemas, Universidad Mayor, Santiago, Chile. ¹⁶⁴Vicerrectoría de Investigación y Postgrado, Universidad de La Frontera, Temuco, Chile. ¹⁶⁵Departamento de Silvicultura y Conservación de la Naturaleza, Universidad de Chile, Temuco, Chile. ¹⁶⁶Rhino and Forest Fund e.V., Kehl, Germany. ¹⁶⁷Siberian Federal University, Krasnoyarsk, Russian Federation. ¹⁶⁸Geobotany, Faculty of Biology, University of Freiburg, Freiburg im Breisgau, Germany. ¹⁶⁹Department of Geography, Remote Sensing Laboratories, University of Zürich, Zurich, Switzerland. ¹⁷⁰National Forest Centre, Forest Research Institute Zvolen, Zvolen, Slovakia. ¹⁷¹Université de Lorraine, AgroParisTech, INRAE, Silva, Nancy, France. ¹⁷²Center for Biodiversity Dynamics in a Changing World (BIOCHANGE), Department of Biology, Aarhus University, Aarhus, Denmark. ¹⁷³Faculty of Environmental Sciences and Natural Resource Management, Norwegian University of Life Sciences, Ås, Norway. ¹⁷⁴Departamento de Biología, Universidad de la Serena, La Serena, Chile. ¹⁷⁵Centro de Ciências Biológicas e da Natureza, Universidade Federal do Acre, Rio Branco, Acre, Brazil. ¹⁷⁶Guyana Forestry Commission, Georgetown, French Guiana. ¹⁷⁷Environmental and Life Sciences, Faculty of Science, Universiti Brunei Darussalam, Gadong, Brunei Darussalam. ¹⁷⁸Plant Systematic and Ecology Laboratory, Department of Biology, Higher Teachers' Training College, University of Yaoundé I, Yaoundé, Cameroon. ¹⁷⁹Departamento de Ecologia, Universidade Federal do Rio Grande do Norte, Natal, Rio Grande do Norte, Brazil. ¹⁸⁰Department of Forest Management, Dendrometry and Forest Economics, Warsaw University of Life Sciences, Warsaw, Poland. ¹⁸¹Center for Ecological Dynamics in a Novel Biosphere (ECONOVO) & Center for Biodiversity Dynamics in a Changing World (BIOCHANGE), Department of Biology, Aarhus University, Aarhus C, Denmark. ¹⁸²Section for Ecoinformatics and Biodiversity, Department of Biology, Aarhus University, Aarhus, Denmark. ¹⁸³Faculty of Forestry and Wood Sciences, Czech University of Life Sciences, Prague, Czech Republic. ¹⁸⁴Wildlife Conservation Society, New York, NY, USA. ¹⁸⁵Quantitative Biodiversity Dynamics, Department of Biology, Utrecht University, Utrecht, the Netherlands. ¹⁸⁶Iwokrama International Centre for Rainforest Conservation and Development (IIC), Georgetown, French Guiana. ¹⁸⁷School of Forestry and Environmental Studies, Yale University, New Haven, CT, USA. ¹⁸⁸Botanical Garden of Ural Branch of Russian Academy of Sciences, Ural State Forest Engineering University, Yekaterinburg, Russian Federation. ¹⁸⁹Pontificia Universidad Católica del Ecuador, Quito, Ecuador. ¹⁹⁰LINCGlobal, Museo Nacional de Ciencias Naturales, CSIC, Madrid, Spain. ¹⁹¹Plant Ecology and Nature Conservation Group, Wageningen University, Wageningen, the Netherlands. ¹⁹²Silviculture Research Institute, Vietnamese Academy of Forest Sciences, Hanoi, Vietnam. ¹⁹³Department of Biology, Stanford University, Stanford, CA, USA. ¹⁹⁴Agricultural High School, ESAV, Polytechnic Institute of Viseu, IPV, Viseu, Portugal. ¹⁹⁵Centre for the Research and Technology of Agro-Environmental and Biological Sciences, CITAB, UTAD, Quinta de Prados, Vila Real, Portugal. ¹⁹⁶Department of Forest Engineering, Universidade Regional de Blumenau, Blumenau, Brazil. ¹⁹⁷Environmental Studies and Research Center, University of Campinas, UNICAMP, Campinas, Brazil. ¹⁹⁸Department of Forest and Wood Science, University of Stellenbosch, Stellenbosch, South Africa. ¹⁹⁹Key Laboratory of Tropical Biological Resources, Ministry of Education, School of Life and Pharmaceutical Sciences, Hainan University, Haikou, China. ²⁰⁰Division of Forestry and Natural Resources, West Virginia University, Morgantown, WV, USA. ²⁰¹Department of Zoology, University of Oxford, Oxford, UK. ²⁰²Manaaki Whenua—Landcare Research, Lincoln, New Zealand. ²⁰³Department of Wetland Ecology, Institute for Geography and Geoecology, Karlsruhe Institute for Technology, Karlsruhe, Germany. ²⁰⁴Independent Researcher, Sommersbergseestrasse, Bad Aussee, Austria. ²⁰⁵Centre for Agricultural Research in Suriname (CELOS), Paramaribo, Suriname. ²⁰⁶Tropenbos International, Wageningen, the Netherlands. ²⁰⁷Polish State Forests, Coordination Center for Environmental Projects, Warsaw, Poland. ²⁰⁸Research Center of Forest Management Engineering of State Forestry and Grassland Administration, Beijing Forestry University, Beijing, China. ²⁰⁹Present address: Department of Genetics, Evolution, and Environment, University College London, London, United Kingdom. e-mail: haozhi.ma@usys.ethz.ch

Reporting Summary

Nature Portfolio wishes to improve the reproducibility of the work that we publish. This form provides structure for consistency and transparency in reporting. For further information on Nature Portfolio policies, see our [Editorial Policies](#) and the [Editorial Policy Checklist](#).

Statistics

For all statistical analyses, confirm that the following items are present in the figure legend, table legend, main text, or Methods section.

n/a Confirmed

- ☒ ☒ The exact sample size (n) for each experimental group/condition, given as a discrete number and unit of measurement
- ☒ ☐ A statement on whether measurements were taken from distinct samples or whether the same sample was measured repeatedly
- ☒ ☐ The statistical test(s) used AND whether they are one- or two-sided
Only common tests should be described solely by name; describe more complex techniques in the Methods section.
- ☐ ☒ A description of all covariates tested
- ☒ ☐ A description of any assumptions or corrections, such as tests of normality and adjustment for multiple comparisons
- ☐ ☒ A full description of the statistical parameters including central tendency (e.g. means) or other basic estimates (e.g. regression coefficient) AND variation (e.g. standard deviation) or associated estimates of uncertainty (e.g. confidence intervals)
- ☒ ☐ For null hypothesis testing, the test statistic (e.g. F , t , r) with confidence intervals, effect sizes, degrees of freedom and P value noted
Give P values as exact values whenever suitable.
- ☒ ☐ For Bayesian analysis, information on the choice of priors and Markov chain Monte Carlo settings
- ☒ ☐ For hierarchical and complex designs, identification of the appropriate level for tests and full reporting of outcomes
- ☒ ☐ Estimates of effect sizes (e.g. Cohen's d , Pearson's r), indicating how they were calculated

Our web collection on [statistics for biologists](#) contains articles on many of the points above.

Software and code

Policy information about [availability of computer code](#)

Data collection No software was used

Data analysis The analyses were run in R (version 4.2.2) [R Core Team (2021) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <http://www.R-project.org>. Accessed 17 Jan 2021] and Google Earth Engine (<https://earthengine.google.com>) (v0.1.361). The code used for this study is available at <https://doi.org/10.5281/zenodo.7967245>

For manuscripts utilizing custom algorithms or software that are central to the research but not yet described in published literature, software must be made available to editors and reviewers. We strongly encourage code deposition in a community repository (e.g. GitHub). See the Nature Portfolio [guidelines for submitting code & software](#) for further information.

Data

Policy information about [availability of data](#)

All manuscripts must include a [data availability statement](#). This statement should provide the following information, where applicable:

- Accession codes, unique identifiers, or web links for publicly available datasets
- A description of any restrictions on data availability
- For clinical datasets or third party data, please ensure that the statement adheres to our [policy](#)

Tree occurrence data is available upon request from the Global Forest Biodiversity initiative (GFBi). Information on leaf habit (evergreen vs. deciduous) and leaf

form (broadleaved vs. needle-leaved) came from the TRY database. The World Soil Information Service (WOSIS) dataset contributes to the model performance evaluation. Additionally, the Tallo dataset is used to test the robustness of the random forest model.

Human research participants

Policy information about [studies involving human research participants and Sex and Gender in Research](#).

Reporting on sex and gender

Population characteristics

Recruitment

Ethics oversight

Note that full information on the approval of the study protocol must also be provided in the manuscript.

Field-specific reporting

Please select the one below that is the best fit for your research. If you are not sure, read the appropriate sections before making your selection.

☐ Life sciences ☐ Behavioural & social sciences ☒ Ecological, evolutionary & environmental sciences

For a reference copy of the document with all sections, see nature.com/documents/nr-reporting-summary-flat.pdf

Ecological, evolutionary & environmental sciences study design

All studies must disclose on these points even when the disclosure is negative.

Study description	We analyzed the global distribution of forest leaf types, integrating ground-sourced information from 9,781 standardized forest inventory plots with leaf habit and leaf form information for a total of 10,274 species from the TRY database. Using random forest models, we quantified the global proportion and distribution of leaf types. Using information on both the occurrences of individuals per plot and the basal-area weighted occurrences of each individual, we model leaf-type proportions at the individual level (individual-based leaf type) and the basal-area level (area-based leaf type). To assess the relative importance of environmental features on variation in leaf types, we ran random forest models including a range of environmental variables. To quantify the proportions of different leaf types across global forests, we combined a global tree density distribution map with our individual-based leaf type models. To estimate biomass stored in each of the leaf types, we incorporate our area-based leaf type models with recently published forest aboveground biomass. Using the documented correlations between forest leaf types and local climate, we project the future leaf type models using three climate-change scenarios (low-emission scenario [SSP1-RCP2.6], business-as-usual scenario [SSP3-RCP7], and high-emission scenario [SSP5-RCP8.5]).
Research sample	Plot-level forest leaf types across the globe.
Sampling strategy	No statistical methods were used to predetermine sample size.
Data collection	Tree occurrence data was from Global Forest Biodiversity initiative (GFBi) dataset. Information on leaf habit (evergreen vs. deciduous) and leaf form (broadleaved vs. needle-leaved) were from TRY plant trait database.
Timing and spatial scale	The average year of observation across all plots was 2005.
Data exclusions	Individuals with stem diameters <10 cm were excluded as the focus was on adult trees, and only plots with ≥10 adult individuals were included in the final analysis. For plots with time series data, only the most recent observation year was included in the analysis.
Reproducibility	n/a
Randomization	n/a
Blinding	n/a
Did the study involve field work?	☐ Yes ☒ No

Reporting for specific materials, systems and methods

We require information from authors about some types of materials, experimental systems and methods used in many studies. Here, indicate whether each material, system or method listed is relevant to your study. If you are not sure if a list item applies to your research, read the appropriate section before selecting a response.

Materials & experimental systems

n/a	Involved in the study
☒	☐ Antibodies
☒	☐ Eukaryotic cell lines
☒	☐ Palaeontology and archaeology
☒	☐ Animals and other organisms
☒	☐ Clinical data
☒	☐ Dual use research of concern

Methods

n/a	Involved in the study
☒	☐ ChIP-seq
☒	☐ Flow cytometry
☒	☐ MRI-based neuroimaging

Terms and Conditions

Springer Nature journal content, brought to you courtesy of Springer Nature Customer Service Center GmbH (“Springer Nature”).

Springer Nature supports a reasonable amount of sharing of research papers by authors, subscribers and authorised users (“Users”), for small-scale personal, non-commercial use provided that all copyright, trade and service marks and other proprietary notices are maintained. By accessing, sharing, receiving or otherwise using the Springer Nature journal content you agree to these terms of use (“Terms”). For these purposes, Springer Nature considers academic use (by researchers and students) to be non-commercial.

These Terms are supplementary and will apply in addition to any applicable website terms and conditions, a relevant site licence or a personal subscription. These Terms will prevail over any conflict or ambiguity with regards to the relevant terms, a site licence or a personal subscription (to the extent of the conflict or ambiguity only). For Creative Commons-licensed articles, the terms of the Creative Commons license used will apply.

We collect and use personal data to provide access to the Springer Nature journal content. We may also use these personal data internally within ResearchGate and Springer Nature and as agreed share it, in an anonymised way, for purposes of tracking, analysis and reporting. We will not otherwise disclose your personal data outside the ResearchGate or the Springer Nature group of companies unless we have your permission as detailed in the Privacy Policy.

While Users may use the Springer Nature journal content for small scale, personal non-commercial use, it is important to note that Users may not:

1. use such content for the purpose of providing other users with access on a regular or large scale basis or as a means to circumvent access control;
2. use such content where to do so would be considered a criminal or statutory offence in any jurisdiction, or gives rise to civil liability, or is otherwise unlawful;
3. falsely or misleadingly imply or suggest endorsement, approval, sponsorship, or association unless explicitly agreed to by Springer Nature in writing;
4. use bots or other automated methods to access the content or redirect messages
5. override any security feature or exclusionary protocol; or
6. share the content in order to create substitute for Springer Nature products or services or a systematic database of Springer Nature journal content.

In line with the restriction against commercial use, Springer Nature does not permit the creation of a product or service that creates revenue, royalties, rent or income from our content or its inclusion as part of a paid for service or for other commercial gain. Springer Nature journal content cannot be used for inter-library loans and librarians may not upload Springer Nature journal content on a large scale into their, or any other, institutional repository.

These terms of use are reviewed regularly and may be amended at any time. Springer Nature is not obligated to publish any information or content on this website and may remove it or features or functionality at our sole discretion, at any time with or without notice. Springer Nature may revoke this licence to you at any time and remove access to any copies of the Springer Nature journal content which have been saved.

To the fullest extent permitted by law, Springer Nature makes no warranties, representations or guarantees to Users, either express or implied with respect to the Springer nature journal content and all parties disclaim and waive any implied warranties or warranties imposed by law, including merchantability or fitness for any particular purpose.

Please note that these rights do not automatically extend to content, data or other material published by Springer Nature that may be licensed from third parties.

If you would like to use or distribute our Springer Nature journal content to a wider audience or on a regular basis or in any other manner not expressly permitted by these Terms, please contact Springer Nature at

onlineservice@springernature.com